



Prevalence of Thyroid Dysfunction Disorders among Adult Populations in the Middle-East: A Systematic Review and Meta-analysis

Shiva Kargar^{1,*}, Seyed Mehdi Tabatabaei¹, Hassan Okati-Aliabad¹ and Hossein Izadi Rad¹

¹Health Promotion Research Center, Zahedan University of Medical Sciences, Zahedan, Iran

Abstract:

Background: Thyroid dysfunction is a systemic disorder that causes severe morbidity and is a public health problem worldwide. This study aimed to evaluate the prevalence of thyroid dysfunction among adults in the Middle East.

Methods and Materials: We searched PubMed, Google Scholar, and Medline databases from 2000-2021 to identify studies that presented the prevalence of thyroid dysfunction, hypothyroidism, subclinical hypothyroidism, hyperthyroidism, and subclinical hyperthyroidism in the Middle East. A random-effects model was used to calculate the pooled prevalence and confidence intervals of thyroid dysfunction. The data were analyzed using STATA-V14.

Results: Generally, 345 studies had eligible criteria to be included in this meta-analysis. The pooled prevalence of thyroid disorders, overt hypothyroidism, subclinical hypothyroidism, overt hyperthyroidism, and subclinical hyperthyroidism in the Middle East were 19.2% (95% CI: 11.0 - 33.2), 7.2% (95% CI: 3.6 - 14.3), 8.3% (95% CI: 5.3 - 13.0), 2.4% (95% CI: 1.4- 3.9), and 3.2% (95% CI: 2.1 - 4.7), respectively. Moreover, the prevalence of thyroid disorders increased from 15.2% (95% CI: 9.8-23.6) to 31.5% (95% CI: 22.5- 44.2) between 2000 and 2022.

Conclusions: Current meta-analysis suggests that thyroid disorders are more prevalent among adults in the Middle East. Moreover, with an increasing trend in the prevalence of thyroid disorders during the last two decades, early screening and prevention of the disease should be practiced.

Keywords: Prevalence, Hyperthyroidism, Hypothyroidism, Thyroid dysfunction, Middle-East, Meta-analysis.

© 2024 The Author(s). Published by Bentham Open.

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International Public License (CC-BY 4.0), a copy of which is available at: <https://creativecommons.org/licenses/by/4.0/legalcode>. This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

*Address correspondence to this author at the Health Promotion Research Center, Zahedan University of Medical Sciences, Zahedan, Iran; E-mail: shivakargar@yahoo.com

Cite as: Kargar S, Tabatabaei S, Okati-Aliabad H, Rad H. Prevalence of Thyroid Dysfunction Disorders among Adult Populations in the Middle-East: A Systematic Review and Meta-analysis. *Open Public Health J*, 2024; 17: e18749445317174. <http://dx.doi.org/10.2174/0118749445317174240827052511>



Received: April 17, 2024
Revised: July 09, 2024
Accepted: July 10, 2024
Published: ?? ??, 2024



Send Orders for Reprints to
reprints@benthamscience.net

1. INTRODUCTION

Thyroid dysfunction is one of the most prevalent endocrine disorders [1]. The thyroid gland generates the thyroid hormones thyroxine (T4) and triiodo-thyronine (T3) in response to pituitary production of thyrotrophin (TSH). Overt hyperthyroidism is characterized by subnormal serum TSH levels and elevated free serum triiodo-thyronine (T3) or free T4 levels. In contrast, subclinical hyperthyroidism is defined as subnormal serum TSH levels and normal free triiodothyronine (FT3) and free thyroid hormone (FT4) or thyroxine levels. On the other hand, Overt hypothyroidism is defined by a decrease in free thyroxine levels (T4) in the

presence of elevated TSH, while Subclinical hypothyroidism occurs when TSH levels are slightly elevated but T3 and T4 or thyroxine levels are normal [2, 3].

Some possible symptoms of hyperthyroidism include exhaustion, weight loss, increased perspiration, heat intolerance, tachycardia, tremor, and hyperactive reflexes that can develop gradually or abruptly [4]. Exhaustion, dry skin, susceptibility to colds, hair loss, weight gain, constipation, changes in voice, and slower movement and thinking are common symptoms of hypothyroidism that, at onset, are typically slow [5].

Thyroid dysfunction is prevalent at different ages, sexes, race/ethnicity, and geographical locations due to differences in dietary iodine intake [6]. Women, older people (> 60 years old), those with a prior personal history of or a significant family history of thyroid illness, and postpartum women are at a higher risk of developing thyroid dysfunction [2].

Overt thyroid dysfunction has been related to different types of cognitive decline, peripheral neuropathy, and psychological health. Moreover, they have effects on the nervous system, cardiovascular system, bone health, and erythropoiesis [6, 7]. Studies have shown that the prevalence of different types of thyroid disease in the Arab world varies from 6.18 to 47.34%, and thyroid cancer is very prevalent in Arab countries. The diet and lifestyle of the Arab countries have changed over the past years, and thus, thyroid diseases are becoming more prevalent, making the economic crisis greater [8].

Limited studies have been conducted on the prevalence of various types of thyroid disease in the Middle East. Therefore, this study aimed to investigate the prevalence of thyroid dysfunction, namely hypothyroidism, and hyperthyroidism, among the adult population in the Middle East.

2. METHODS

2.1. Search Strategy and Selection Criteria

This meta-analysis was conducted according to the Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA) guidelines [9]. A literature search was conducted in PubMed, Google Scholar, and Medline databases to search for studies that presented the prevalence of thyroid dysfunction, hypothyroidism, subclinical hypothyroidism, hyperthyroidism, and subclinical hyperthyroidism in the Middle East. The search was carried out using the search terms “prevalence”, “hyperthyroidism”, “hypothyroidism”, “subclinical hyperthyroidism”, “subclinical hypothyroidism”, “Thyroid”,

“Thyroid Dysfunction Disorders”, “The Middle East”, and “countries in the Middle East area”. The search was limited to studies that were published between 2000 and July 2021.

2.2. Inclusion and Exclusion Criteria

Studies were included if they provided sufficient information to estimate the prevalence with confidence intervals of thyroid dysfunction, which were cross-sectional studies published in the English language. The exclusion criteria were review, not providing sufficient information, not cross-sectional studies, involving adolescents and children, focusing on specific groups like patients with diabetes, hypertension, postmenopausal women, pregnant women, etc. Moreover, the studies measuring the prevalence of Congenital hypothyroidism and Postpartum Thyroiditis were excluded as well.

2.3. Study Selection and Data Extraction

At first, we screened all of the identified studies through databases based on titles and abstracts. Then, the full text of relevant articles was read, and data extraction was performed from eligible studies. The data extracted from the studies included names of authors, year of publication, setting (country’s name, city), sampling method, diagnostic criteria, gender, number of participants, age, prevalence of Thyroid Dysfunction, and its 95% confidence interval. Table 1 illustrates the main characteristics of the included studies in this meta-analysis.

2.4. Statistical Analysis

The data were analyzed using STATA-14 (Stata Corp, Texas, USA) statistical software. Pooled estimates of thyroid dysfunction prevalence and confidence intervals were calculated using a random-effects model. The I2 test was used to measure study heterogeneity. Subgroup analysis was then performed to investigate the sources of heterogeneity. Egger’s test was also performed to check for publication bias.

Table 1. The characteristics of the included studies.

Authors/ Year of Publication/Refs	Setting		Sampling				Dyslipidaemias		Overt Hypothyroidism *		Subclinical Hypothyroidism **		Overt Hyperthyroidism †		Subclinical Hyperthyroidism ‡	
	Country	City	Age	Method	Sex	N	Criteria	Prevalence (95% CI)	*Criteria	Prevalence (95% CI)	Criteria	Prevalence (95% CI)	Criteria	Prevalence (95% CI)	Criteria	Prevalence (95% CI)
Elham Faghih Imani,2010 [10]	Iran	isfahan	> 20	multistage clustering	FM	263	-	-	5	6.5 (3.8 - 10.1)	1	6.9 (4.1- 10.6)	1	0.3 (0.01 - 2.1)	1	0.7 (0.09 - 2.7)
-	-	-	-	-	M	142	-	-	-	3.5 (1.1 - 8)	-	5.6(2.4 - 10.8)	-	-	-	0.7 (0.02 - 3.8)
-	-	-	-	-	F	121	-	-	-	10.7 (5.8 - 17.6)	-	7.4(3.4 - 13.6)	-	0.8 (0.02 - 4.5)	-	0.8 (0.02 - 4.5)
Ebrahim Barzegari,2021 [11]	Iran	Ravansar	35-65	-	FM	10069	-	-	-	3.21 (2.8 - 3.5)	-	-	-	0.44 (0.3 - 0.5)	-	-
-	-	-	-	-	M	4752	-	-	-	0.88 (0.6 - 1.1)	-	-	-	0.35(0.21- 0.57)	-	-
-	-	-	-	-	F	5269	-	-	-	5.36 (4.7 - 5.9)	-	-	-	0.51(0.34- 0.75)	-	-
Bahram Eshraghi,2022 [12]	Iran	Rasht	39.5±13.45	-	FM	383	-	-	-	8.9(6 - 11.8)	-	-	-	89 (85.4- 91.9)	-	-
-	-	-	-	-	M	155	-	-	-	3.8 (1.4- 8.2)	-	-	-	94.8 (90 - 97.7)	-	-
-	-	-	-	-	F	228	-	-	-	11.8 (7.9 - 16.7)	-	-	-	85(79.7 - 89.4)	-	-

Authors/ Year of Publication/Refs	Setting		Sampling				Dyslipidaemias		Overt Hypothyroidism *		Subclinical Hypothyroidism **		Overt Hyperthyroidism †		Subclinical Hyperthyroidism ‡	
	Country	City	Age	Method	Sex	N	Criteria	Prevalence (95% CI)	*Criteria	Prevalence (95% CI)	Criteria	Prevalence (95% CI)	Criteria	Prevalence (95% CI)	Criteria	Prevalence (95% CI)
NIAFAR M,2009 [13]	Iran	East Azerbaijan	60-89	random	FM	1000	-	12.7(10.7 - 14.9)	-	1.5(0.8 - 2.4)	-	5.8 (4.4 - 7.4)	-	1.2 (0.6 - 2.09)	-	4.1 (2.9 - 5.5)
Maryam Rezaei,2019 [14]	Iran	Birjand	-	voluntarily	FM	110	-	-	11	30 (21.6 - 39.4)	-	-	1	30 (21.6 - 39.4)	-	-
-	-	-	-	-	M	45	-	-	-	22.2 (11.2 - 37)	-	-	-	22.2 (11.2 - 37)	-	-
-	-	-	-	-	F	65	-	-	-	35.3(23.9-48.2)	-	-	-	35.3(23.9-48.2)	-	-
Ladan Mehran,2017 [15]	Iran	Tehran	40.3 ± 14.4	multistage cluster random	FM	5422	-	-	1	1.8(1.5 - 2.2)	1	5.4(4.8 - 6.06)	1	1.5(1.2 - 1.8)	1	3.2 (2.8 - 3.7)
-	-	-	-	-	M	2318	-	-	-	16.8(15.3 - 18.4)	-	23.7 (22 - 25.5)	-	42.7 (40.6 - 44.7)	-	40.4(38.3 - 42.4)
-	-	-	-	-	F	3104	-	-	-	83.2 (81.8 - 84.4)	-	76.2(74.6 - 77.6)	-	57.2(55.5 - 59)	-	59.6 (57.8 - 61.3)
Ashraf Aminorroaya,2008 [16]	Iran	Isfahan	> 20	multistage cluster	FM	2523	-	-	-	-	-	-	1	0.8(0.5 - 1.2)	1	0.9(0.6 - 1.4)
-	-	-	-	-	M	1275	-	-	-	-	-	-	-	0.3 (0.1 - 0.9)	-	0.7 (0.3 - 1.3)
-	-	-	-	-	F	1248	-	-	-	-	-	-	-	1.2 (0.7 - 2.07)	-	1.2 (0.7 - 2.07)
Farnaz Rahmani,2018 [17]	Iran	Tabriz	18-58	convenient sampling	FM	261	-	-	-	1.1(0.2 - 3.3)	-	16.8(12.5- 21.9)	-	0.3 (0.01 - 2.1)	-	-
Yusuf Aydin,2014 [18]	Turkey	-	18-92	-	FM	2233	-	-	1	6.5 (5.5 - 7.6)	1	12 (10.6 - 13.4)	-	0.5 (0.2 - 0.9)	-	11 (9.7 - 12.4)
-	-	-	-	-	M	803	-	-	-	1.7 (0.9 - 2.9)	-	2.8(1.8 - 4.2)	-	0.6 (0.2 - 1.4)	-	10.8 (8.7 - 13.1)
-	-	-	-	-	F	1430	-	-	-	1.3(0.8 - 2)	-	4.4 (3.4 - 5.6)	-	0.4 (0.2 - 1)	-	11.1 (9.6 - 12.9)
Mustafa Behçet Demirbaş,2019 [19]	Turkey	Istanbul	> 65	-	FM	500	-	-	-	0.6 (0.1 - 1.7)	-	2.8 (1.5- 4.6)	-	0.6 (0.1-1.7)	-	16 (12.9 - 19.5)
Faruk Kutluturk,2013 [20]	Turkey	province of Tokat	>18	random sampling	FM	1095	-	-	2	1.6 (1 - 2.6)	2	2.7 (1.9 - 4)	1	0.5 (0.2 - 1.2)	1	4.9 (3.8 - 6.6)
-	-	-	-	-	M	541	-	-	-	0.7 (0.2 - 1.8)	-	1.8 (0.8 - 3.3)	-	0.4 (0.05 - 1.3)	-	3.5(2.1 - 5.4)
-	-	-	-	-	F	554	-	-	-	2.5(1.4 - 4.2)	-	3.6 (2.2 - 5.6)	-	0.7 (0.2 - 1.8)	-	6.4 (4.5 - 8.8)
Glay HERGEN,2005 [21]	Turkey	-	>34	-	FM	512	-	-	3	3.1 (1.8 - 5)	-	-	1	5.8 (3.9 - 8.2)	-	-
-	-	-	-	-	M	239	-	-	-	1.7(0.4 - 4.2)	-	-	-	7 (4.2 - 11.1)	-	-
-	-	-	-	-	F	273	-	-	-	4.4(2.2 - 7.5)	-	-	-	5 (2.8 - 8.4)	-	-
Selahittin Çayan,2017 [22]	Turkey	19 provinces of Turkey	≥40	randomly	FM	2760	-	-	-	0.5 (0.2 - 0.8)	-	-	-	1.6 (1.1 - 2.1)	-	-
Ayktut Sarıtaş,2015 [23]	Turkey	Giresun	≥18	-	FM	10600	-	-	4	1.9(1.6 - 2.1)	1	6.7 (6.2 - 7.1)	3	0.7 (0.5 - 0.8)	2	1.8 (1.5 - 2)
-	-	-	-	-	M	5697	-	-	-	1.4(1.1 - 1.7)	-	5.2 (4.6 - 5.8)	-	0.6 (0.4 - 0.8)	-	1.7 (1.3 - 2)
-	-	-	-	-	F	4903	-	-	-	2.4 (2 - 2.9)	-	8.3 (7.5 - 9.1)	-	0.6 (0.4 - 0.9)	-	1.9 (1.6 - 2.4)
Rana Hasanato,2017 [24]	Saudi Arabia	-	20-65	-	F	199	-	-	2	10(6.2 - 15)	-	-	-	3 (1.1 - 6.4)	-	-
Ahmed Ali Gaffer Ali,2016 [25]	Saudi Arabia	AlBAHAH city	≥18	-	FM	71	-	43.6(31.9 - 55.9)	-	40.8(29.3 - 53.1)	-	-	-	2.8 (0.03 - 9.8)	-	-
-	-	-	-	-	M	8	-	25 (3.1 - 65)	-	25 (3.1 - 65)	-	-	-	-	-	-
-	-	-	-	-	F	63	-	46.03(33.3 - 59)	-	42.8 (30.4 - 55.9)	-	-	-	3.17(0.3 - 11)	-	-
18Malak A. Al-Shammari,2022 [26]	Saudi Arabia	Dammam City	median: 37	-	FM	240	-	-	5	29.2 (23.5- 35.3)	-	-	4	15 (10.7 - 20.1)	-	-
-	-	-	-	-	M	35	-	-	-	-	-	-	-	-	-	-
-	-	-	-	-	F	205	-	-	-	-	-	-	-	-	-	-
19 Saif Aboud M. Alqahtani,2021 [27]	Saudi Arabia	Asir region	43.4 ± 15.8	-	FM	9992	-	49.7(48.7- 50.7)	6	5.3(4.8 - 5.7)	1	39.2 (38.2 - 40.2)	1	2.49 (2.2 - 2.8)	1	2.72(2.4 - 3)
-	-	-	-	-	M	-	-	18.3(17.6- 19.1)	-	2.4(2.1 - 2.7)	-	15.3 (14.3- 15.7)	-	1.1 (0.9 - 1.3)	-	0.8(0.7 - 1)
-	-	-	-	-	F	-	-	31.9(30.9- 32.8)	-	2.9(2.5 - 3.2)	-	24.2(23.3- 25)	-	1.4 (1.1 - 1.6)	-	1.8(1.5 - 2.1)
20 Shahad Lafi Alanazi,2018 [28]	Saudi Arabia	Arar	>20	multistage stratified random	FM	160	-	22.5(16.2- 29.7)	7	15.6 (10.3 - 22.2)	-	-	-	6.8 (3.4 - 11.9)	-	-
-	-	-	-	-	M	-	-	17.3(4.9- 38.7)	-	-	-	-	-	39.1(19.7 - 61.4)	-	-
-	-	-	-	-	F	-	-	15.3(9.7- 22.4)	-	-	-	-	-	1.4 (0.1 - 5.1)	-	-
21 Atheer Mohammed D Alotaibi,2018 [29]	Saudi Arabia	Riyadh city	>18	Systematic random	FM	870	-	-	8	16.7(14.3 - 19.4)	-	-	5	2 (1.1 - 3.1)	-	-

(Table 3) contd....

Authors/ Year of Publication/Refs	Setting		Sampling				Dyslipidaemias		Overt Hypothyroidism *		Subclinical Hypothyroidism **		Overt Hyperthyroidism †		Subclinical Hyperthyroidism ‡	
	Country	City	Age	Method	Sex	N	Criteria	Prevalence (95% CI)	*Criteria	Prevalence (95% CI)	Criteria	Prevalence (95% CI)	Criteria	Prevalence (95% CI)	Criteria	Prevalence (95% CI)
-	-	-	-	-	M	297	-	-	-	-	-	-	-	-	-	-
-	-	-	-	-	F	573	-	-	-	-	-	-	-	-	-	-
24 Eidan Al Eidan,2018 [30]	Saudi Arabia	Riyadh	>18	-	FM	394	-	-	-	1	10.3(7.2- 14)	-	-	1	2.1(0.8 - 4.2)	
25 Bassem Refaat,2015 [31]	Saudi Arabia	Makkah	18-45	-	F	600	-	19.6(16.5- 23)	1	6.3(4.5 - 8.5)	1	-	6	1.6(0.8 - 3)	-	
26 Abdelhameed A Fureeh,2019 [32]	Saudi Arabia	l-Baha	>18	-	FM	567	-	-	2	-	1	15.9 (12.9 -19.1)	1	-	3	4.4(2.8 - 6.4)
-	-	-	-	-	M	101	-	-	-	-	-	2.6 (1.4- 4.3)	-	-	-	0.7(0.1 - 1.7)
-	-	-	-	-	F	466	-	-	-	-	-	13.7(10.9- 16.9)	-	-	-	3.7(2.4 - 5.8)
27 Mohammed Qashgry,2020 [33]	Saudi Arabia	Jeddah	16-56	-	FM	346	-	-	-	-	-	-	-	1.1(0.3 -2.9)	-	-
-	-	-	-	-	M	152	-	-	-	-	-	-	-	-	-	-
-	-	-	-	-	F	194	-	-	-	-	-	-	-	-	-	-
34 Abdulaziz A Alghaihy,2013 [34]	Saudi Arabia	Almadinah Almounawarah	>14	-	FM	177	-	-	-	34.4(27.4 - 41.9)	-	19.2(13.6 -25.7)	-	12.4 (7.9 - 18.2)	-	16.3(11.2 - 22.6)
-	-	-	-	-	M	34	-	-	-	23.5(10.7 41.1)	-	14.7 (4.9- 31)	-	14.7 (4.9- 31)	-	11.7(3.3 - 27.4)
-	-	-	-	-	F	143	-	-	-	37(29.1 - 45.5)	-	20.2(14 - 27.8)	-	11.8(7 - 18.3)	-	17.4(11.6 - 24.7)
35 Abdulwahab Alyahya,2021 [35]	Saudi Arabia	Al Ahsa	18-60	randomly	FM	882	-	-	-	11.7(9.7- 14.1)	-	-	-	1.7(0.2 - 2.7)	-	-
-	-	-	-	-	M	391	-	-	-	-	-	-	-	-	-	-
-	-	-	-	-	F	491	-	-	-	-	-	-	-	-	-	-
36 Khalid SJ Aljabri,2019 [36]	Saudi Arabia	Jeddah	12-105	-	FM	3872	-	-	5	29.1(27.6 -30.5)	-	-	-	-	-	-
-	-	-	-	-	M	884	-	-	-	18.2(15.7- 20.9)	-	-	-	-	-	-
-	-	-	-	-	F	2988	-	-	-	32.2(30.5- 33.9)	-	-	-	-	-	-
38 Nida Suhail,2020 [37]	Saudi Arabia	Arar	>18	-	FM	150	-	-	-	48.6(40.4 - 56.9)	-	-	-	12.6(7.8 - 19)	-	-
-	-	-	-	-	M	64	-	-	-	45.3(32.8 - 58.2)	-	-	-	29.6(18.9 - 42.4)	-	-
-	-	-	-	-	F	86	-	-	-	51.1(40.1 - 62.1)	-	-	-	13.9(7.4 - 23.1)	-	-
28 Munir Abu- Helalah,2020 [38]	Jordan	Karak	18-79	-	FM	7085	-	-	2	14.8(14 - 15.7)	1	5.5 (5 - 6)	1	1.9(1.6 - 2.2)	4	-
-	-	-	-	-	M	2007	-	-	-	9.1(7.8- 10.4)	-	4.3 (3- 5.3)	-	2.2(1.6 - 2.9)	-	-
-	-	-	-	-	F	5084	-	-	-	17.2(16.1- 18.2)	-	5.9(5.3 - 6.6)	-	1.8(1.4 - 2.1)	-	-
29 Burhan Abdullah Zaman,2021 [39]	Iraq	Duhok city	-	-	FM	24568	-	-	-	1.2(1.07 - 1.34)	-	94.8(94.5- 95.1)	-	0.31(0.25 - 0.39)	4	2.2(2.02 - 2.39)
-	-	-	-	-	M	8063	-	-	-	0.9 (0.7 - 1.1)	-	95.1(94.6- 95.6)	-	0.3(0.2 -0.4)	-	2.2(1.8 - 2.5)
-	-	-	-	-	F	16505	-	-	-	1.3(1.1 - 1.5)	-	94.7 (94.3 -95)	-	0.3(0.2 - 0.4)	-	2.2(1.9 - 2.4)
30 Ahmed M Athab Al-Msari,2014 [40]	Iraq	Baquba City	-	-	FM	2973	-	-	-	7.1 (6.2 - 8.1)	-	-	-	5(4.2 - 5.8)	-	-
-	-	-	-	-	M	647	-	-	-	3.5(2.2 - 5.2)	-	-	-	3.7(2.3 - 5.4)	-	-
-	-	-	-	-	F	2326	-	-	-	8.1(7 - 9.3)	-	-	-	5.3(4.4 - 6.3)	-	-
31 Sabah Muhammed Salih,2021 [41]	Iraq	Kirkuk	20-50	-	FM	88	-	-	9	22.7 (14.4- 32.8)	-	-	1	23.8(15.4 - 34.1)	-	-
-	-	-	-	-	M	20	-	-	-	5(0.1 - 24.8)	-	-	-	20(5.7 - 43.6)	-	-
-	-	-	-	-	F	68	-	-	-	27.9 (17.7 - 40.1)	-	-	-	25(15.2 - 36.9)	-	-
32 Payman A. Hamasaheed,2019 [42]	Iraq	Erbil City	-	-	F	433	-	-	-	3(1.6 - 5)	-	-	1	9.4(6.8 -12.6)	-	-
33 Fadhluddin Nasruddin Shakor,2019 [43]	Iraq	Sulaimaniyah city	-	-	FM	115	-	-	2	35.6(26.9 - 45.1)	-	-	-	44.3(35 -53.9)	-	-
-	-	-	-	-	M	14	-	-	-	28.5(8.3- 58.1)	-	-	-	64.2(35.1 - 87.2)	-	-
-	-	-	-	-	F	101	-	-	-	36.6(27.2 - 46.8)	-	-	-	41.5(31.8 - 51.8)	-	-
37 Noor Thair Tahir,2021 [44]	Iraq	Baghdad	12- 62	-	FM	1800	-	-	10	3.2(2.4 - 4.1)	1	14.1(12.5- 15.8)	1	3(2.2 -3.9)	3	4 (3 - 4.9)
-	-	-	-	-	M	388	-	-	-	3.3(1.8 - 5.6)	-	12.8(9.7 - 16.6)	-	3.8(2.1 -6.3)	-	3.3 (1.8 - 5.6)
-	-	-	-	-	F	1412	-	-	-	3.1(2.3 - 4.2)	-	14.4(12.6 - 16.3)	-	2.7(1.9 - 3.7)	-	4.1 (3.1 - 5.2)
39 Hamed Abdel- Aziz Deraz,2019 [45]	Egypt	Zagazig city	67.6± 7.1	-	FM	126	-	13.3(8.3-19.8)	-	1.3(0.1 - 4.7)	-	7.3(3.7 - 12.7)	-	1.3 (0.1- 4.7)	-	3.3 (1 - 7.6)

(Table 3) contd....

Authors/ Year of Publication/Refs	Setting		Sampling				Dyslipidaemias		Overt Hypothyroidism *		Subclinical Hypothyroidism **		Overt Hyperthyroidism †		Subclinical Hyperthyroidism ‡	
	Country	City	Age	Method	Sex	N	Criteria	Prevalence (95% CI)	*Criteria	Prevalence (95% CI)	Criteria	Prevalence (95% CI)	Criteria	Prevalence (95% CI)	Criteria	Prevalence (95% CI)
-	-	-	-	-	M	41	-	19.5(8.8-34.8)	-	-	-	7.3 (1.5 - 19.9)	-	4.8(0.6 -16.5)	-	7.3 (1.5 - 19.9)
-	-	-	-	-	F	85	-	14.1(7.5-23.3)	-	2.3(0.2 - 8.2)	-	9.4 (4.1 - 17.7)	-	-	-	2.3 (0.2 - 8.2)
Amira H. Mahmoud,2005 [46]	Egypt	Qatif city	>60	-	FM	100	-	20(12.6 -29.1)	2	10(4.9 - 17.6)	1	5 (1.6 - 11.2)	1	3(0.6 -8.5)	4	2 (0.2 - 7)
22 Nearemeen M. Rashad,2019 [47]	Egypt	-	-	-	FM	430	-	29.3(25 -33.8)	5	20.4(16.7 - 24.5)	1	44.4(39.6 - 49.2)	4	19.2(15.4 - 23.1)	4	15.8(12.4 - 19.6)

Note: * Overt Hypothyroidism: 1= TSH > 5.06 and FT4 < 0.91 µIU/ml / 2= high TSH level with normal or below normal T4 levels / 3= TSH> 4.2 µU/ml / 4= FT4 <0.7 ng dL-1, (TSH) >4 mIU mL-1 / 5= FT3 < 3.71 pg/ml or FT4 < 1.48 ng/dl or TSH > 4.94 µIU/ml / 6= TSH > 4.94 IU/mL and FT4 < 0.7 ng/dL / 7= symptoms of thyroid dysfunction./ 8= Self report/ 9=TSH >7 µIU/ml / 10=TSH>5.0 (µmol/L), T4<60 (nmol/L) / 11=TSH > 6.3 (mIU/L), T4 (> 12.9 µg/dL), T3 (> 220 ng/dL), FT3 (> 4.2 pg/mL), and FT4 (> 2.0 ng/mL).

** Subclinical Hypothyroidism: 1= TSH > 4 mIU/L and T4, T3, and FT4I were within the normal range/ 2= (TSH: 1, and FT4:N)/

Overt Hyperthyroidism: 1= low TSH (<0.3 mIU/L)with high FT4I or T3 / 3= FT4 >1.7 ng dL-1 and TSH <0.1 mIU mL-1/ 4=FT3 > 3.71 pg/ml or FT4> 1.48 ng/dl or TSH < 4.94 µIU/ml / 5= Selfe report/ 6= TSH < 0.10 µIU/ml and FT4 >22 pmol/L/

\$ Subclinical Hyperthyroidism: 1= Low TSH(<0.3 mIU/L) with normal FT4I and T3/ 2= TSH <0.1 mIU mL-1 with normal free hormone levels/ 3= TSH <0.2 with normal FT4/ 4= Low TSH and normal T3 and T4.




Fig. (1). Flowchart of the present study selection.

3. RESULT

A total of 834 articles were identified from a database search, of which 356 articles were removed due to duplication. Next, 478 articles that were not duplicated were screened, and 281 articles were excluded based on irrelevant titles and abstracts. Then, 197 articles were selected to assess the full text. Of these, 159 articles were excluded for the following reasons: absence of prevalence rate, not providing sufficient information to calculate CI 95%, not finding the full-text article, not being cross-sectional, and being carried out among special groups. Eventually, 38 articles from Iran, Saudi Arabia, Egypt, Iraq, Turkey, and Jordan were eligible to be included in the meta-analysis (Fig. 1). We could not find eligible articles from other countries in the Middle East region.

Four thyroid disorders were described in this survey. The number of studies on the four thyroid disorders was 34 on overt hypothyroidism, 18 on subclinical hypothyroidism, 35 on overt hyperthyroidism, and 17 on subclinical hyperthyroidism.

The pooled prevalence of thyroid disorders was 19.2% (95% CI: 11.0 - 33.2) in the Middle East. The highest prevalence of thyroid disorders was observed in Saudi Arabia at 31.3% (95% CI: 17.9 - 54.9), and the lowest prevalence was in Iran at 12.7% (95% CI: 10.7 - 14.9). Some heterogeneity was seen among the study's findings ($p < 0.001$). In addition, the pooled prevalence of overt hypothyroidism, subclinical hypothyroidism, overt hyperthyroidism, and subclinical hyperthyroidism in the Middle East was 7.2% (95% CI: 3.6 - 14.3), 8.3% (95% CI: 5.3 - 13.0), 2.4% (95% CI: 1.4- 3.9), and 3.2% (95% CI: 2.1 - 4.7), respectively (Table 2).

Although analysis of the data by sex showed that the prevalence of thyroid disorders, overt hypothyroidism, subclinical hypothyroidism, and subclinical hyperthyroidism was higher in women than in men, they were not statistically significant ($p > 0.5$).

Analysis of the data by year of publication of articles illustrated an increasing trend in the prevalence of thyroid disorders between 2000 and 2022, which increased from 15.2% (95% CI: 9.8-23.6) during 2000 - 2010 to 31.5% (95% CI: 22.5- 44.2) during 2016 - 2022, overall ($P < 0.001$) (Figs. 2-4).

4. DISCUSSION

This study investigated the prevalence of thyroid disorders, including hypothyroidism and hyperthyroidism, in the Middle East region. The prevalence of thyroid disorders was high in this region [19.2% (95% CI: 11.0 - 33.2)]. Besides, the frequency of other non-communicable diseases, including hypertension, obesity, etc., was found to be considerable among the adult population in the Middle East [48, 49], which could explain the high prevalence of thyroid disorders.

Moreover, the overall prevalence of thyroid disorders was different among countries, ranging from 31.3 in Saudi Arabia to 12.7 in Iran. The high prevalence of thyroid disorders in Saudi Arabia can be attributed to the reasons mentioned in previous studies, including iodine deficiency and improper nutrition [8]. However, the differences observed in the prevalence of different thyroid lesions among countries in the Middle East may be due to diversity in the study methods, environmental factors, iodination status, and different cut-off values to identify thyroid disorders.

Table 2. The prevalence of thyroid dysfunction disorders in middle east countries.

Variables	N. of Studies (population)	Thyroid Dysfunction	Test for Heterogeneity	Overt Hypothyroidism	Test for Heterogeneity	Subclinical Hypothyroidism	Test for Heterogeneity	Overt Hyperthyroidism	Test for Heterogeneity	Subclinical Hyperthyroidism	Test for Heterogeneity
		Prevalence (95% CI)	(P- value)	Prevalence (95% CI)	(P- value)	Prevalence (95% CI)	(P- value)	Prevalence (95% CI)	(P- value)	Prevalence (95% CI)	(P- value)
Country	-	-	-	-	-	-	-	-	-	-	-
Saudi Arabia	14 (18520)	31.3 (17.9 - 54.9)	P < 0.001	17.9 (10.5 - 30.3)	P < 0.001	18.9 (9.6 - 37.1)	P < 0.001	4.0 (2.1 - 7.6)	P < 0.001	4.6 (1.7 - 11.8)	P < 0.001
Turkey	6 (17700)	-		1.7 (0.8 - 3.6)		5.2 (3.2 - 8.6)		1.05 (0.4 - 2.4)		6.2 (2.1 - 17.9)	
Iran	8 (20031)	12.7 (10.7 - 14.9)		4.2 (2.04 - 9.0)		7.7 (4.5 - 13.09)		2.05 (0.2 - 15.5)		2.05 (1.06 - 3.9)	
Jordan	1 (7085)	-		14.4 (14.0 - 15.7)		5.5 (5.0 - 6.0)		1.9 (1.6 - 2.2)		-	
Egypt	3 (656)	20.4 (12.5 - 33.3)		9.9 (3.8 - 25.9)		12.2 (2.6 - 57.7)		5.02 (0.8 - 29.2)		5.5 (1.3 - 22.3)	
Iraq	6 (29977)	-		6.3 (2.1 - 19.2)		36.5 (5.6 - 236.8)		5.9 (1.3 - 25.6)		2.9 (1.6 - 5.2)	
Total (Middle-East)	38 (93969)	19.2 (11.0 - 33.2)		7.2 (3.6 - 14.3)		8.3 (5.3 - 13.0)		2.4 (1.4 - 3.9)		3.2 (2.1 - 4.7)	
sex	-	-	-	-	-	-	-	-	-	-	-
Female	28 (54298)	26.5 (18.4 - 38.04)	P = 0.5	8.9 (5.1 - 15.4)	P = 0.2	14.1 (9.6 - 20.8)	P = 0.2	3.6 (1.9 - 6.6)	P = 0.6	4.2 (1.3 - 13.7)	P = 0.7
Male	24 (29774)	18.3 (17.5 - 19.07)		5.4 (3.1 - 9.5)		8.1 (3.6 - 18.2)		4.6 (2.3 - 9.07)		3.07 (0.9 - 10.07)	
Year of publication	-	-	-	-	-	-	-	-	-	-	-
2000 - 2010	4 (4298)	15.2 (9.8 - 23.6)	P = 0.01	4.1 (1.8 - 9.0)	P = 0.3	5.9 (4.7 - 7.4)	P = 0.02	1.6 (0.5 - 4.9)	P = 0.4	1.6 (0.5 - 4.8)	P = 0.2
2011 - 2015	6 (17678)	19.6 (16.5 - 23.0)		5.6 (2.4 - 12.8)		8.1 (4.9 - 13.4)		1.6 (0.5 - 4.8)		6.2 (2.1 - 18.4)	
2016 - 2022	28 (71993)	31.5 (22.5 - 44.2)		8.0 (5.0 - 12.6)		13.2 (7.7 - 22.6)		3.7 (1.4 - 9.8)		4.3 (2.6 - 7.2)	




Fig. (2). The prevalence of overt hypothyroidism and subclinical hypothyroidism in the Middle East.




Fig. (3). The prevalence of overt hyperthyroidism and subclinical hyperthyroidism in the Middle East.

The pooled prevalence of overt hypothyroidism, subclinical hypothyroidism, overt hyperthyroidism, and subclinical hyperthyroidism in the Middle East was 7.2% (95% CI: 3.6 - 14.3), 8.3% (95% CI: 5.3 - 13.0), 2.4% (95% CI: 1.4- 3.9), and 3.2% (95% CI: 2.1 - 4.7), respectively. A previous meta-analysis on thyroid dysfunctions estimated the prevalence of undiagnosed hyperthyroidism in Europe at 1.7% (95% CI: 1.66-1.88), with a clear predominance of the subclinical form of the disease: overt hyperthyroidism 0.35% (95% CI: 0.29-0.41) and subclinical hyperthyroidism 0.50% (95% CI: 0.57-1.43) [50]. Moreover, overt hypothyroidism and subclinical hypothyroidism were

found in 0.65% (95% CI 0.38-0.99) and 4.11% (95% CI 3.05-5.31) of Europe, respectively [51]. Comparing these results with the present meta-analysis results, the Middle East region exceeds Europe by 6.5% and 2% for overt hypothyroidism and hyperthyroidism and by 4.1% and 2.7% for subclinical hypothyroidism and hyperthyroidism, respectively. Moreover, the prevalence of hypothyroidism and hyperthyroidism among adults in the United States has been calculated at 9.47% and 1.19%, respectively [52]. Therefore, the prevalence of hypothyroidism and hyperthyroidism are roughly similar in the Middle East and the United States (0.7% versus 9.47%, 2.4% versus 1.9%).




Fig. (4). Trend and prevalence of thyroid disorders, overt hypothyroidism, subclinical hypothyroidism, overt hyperthyroidism, and subclinical hyperthyroidism from 2000 to 2022.

Non-communicable diseases have increased among adults in the Middle East during the last two decades. Moreover, the findings of this meta-analysis show an increasing trend in the prevalence of thyroid disorders between 2000 and 2022. Changing lifestyles, using an unhealthy diet (high-calorie and fat), obesity, using alcohol, and smoking in this regain could be possible reasons for an increasing trend in the prevalence of thyroid disorders [11, 48, 53].

The present results, like other studies [51], point out that the prevalence of subclinical hypothyroidism (8.3%) and subclinical hyperthyroidism (3.2%) is higher than overt hypothyroidism (7.2%) and overt hyperthyroidism (2.4%). This high prevalence sheds light on the potential for the underappreciation of subclinical thyroid diseases, which can have significant effects on the management and treatment of adult patients. Our findings suggested that additional thyroid function screening may be necessary among adults, as well as increased awareness of thyroid dysfunction.

The prevalence of thyroid dysfunction varies by age, sex, race/ethnicity, and iodine status of populations [54]. In this study, the prevalence of thyroid disorders, overt hypothyroidism, subclinical hypothyroidism, and subclinical hyperthyroidism was higher in women compared with men. Moreover, previous studies have shown that the prevalence of hypothyroidism has been higher in females [6, 30, 51]. The strongest support for screening comes from the 2005 AACE, American Thyroid Association (ATA), and Endocrine Society Consensus Statement, which

suggested thyroid screening for all women starting at age 35 years and continuing every five years after that [50].

CONCLUSION

According to the results of the current meta-analysis, the prevalence of thyroid disorders was high among adults, especially women in the Middle East. Moreover, an increasing trend was observed in the prevalence of thyroid disorders during the last two decades. Therefore, this meta-analysis suggests the need for increased screening of thyroid disorders and prevention of disease at the primary level by educating the population.

LIMITATIONS

one of the limitations of this meta-analysis was that the cut-off value of serum thyrotropin (TSH), thyroxine (T4), and triiodothyronine (T3) to determine the four thyroid disorders was not identical between the studies. In addition, the years of publication of articles and the distribution of age were different between studies from different countries.

AUTHORS' CONTRIBUTIONS

It is hereby acknowledged that all authors have accepted responsibility for the manuscript's content and consented to its submission. They have meticulously reviewed all results and unanimously approved the final version of the manuscript.

LIST OF ABBREVIATIONS

TSH	= Serum Thyrotropin
T4	= Thyroxine
T3	= Triiodothyronine
PRISMA	= Preferred Reporting Items for Systematic Reviews and Meta-Analysis
ATA	= American Thyroid Association

CONSENT FOR PUBLICATION

Not applicable.

STANDARDS OF REPORTING

PRISMA guidelines and methodologies were followed.

AVAILABILITY OF DATA AND MATERIAL

All the data and supportive information are provided within the article.

FUNDING

None.

CONFLICT OF INTEREST

The authors declare no conflict of interest, financial or otherwise.

ACKNOWLEDGEMENTS

Declared none.

SUPPLEMENTARY MATERIAL

PRISMA checklist is available as supplementary material on the publisher's website along with the published article.

REFERENCES

- [1] Kalra S, Aggarwal S, Khandelwal D. Thyroid dysfunction and dysmetabolic syndrome: The need for enhanced thyrovigilance strategies. *Int J Endocrinol* 2021; 2021: 9641846. <http://dx.doi.org/10.1155/2021/9641846>
- [2] Reyes Domingo F, Avey MT, Doull M. Screening for thyroid dysfunction and treatment of screen-detected thyroid dysfunction in asymptomatic, community-dwelling adults: A systematic review. *Syst Rev* 2019; 8(1): 260. <http://dx.doi.org/10.1186/s13643-019-1181-7> PMID: 31735166
- [3] Floriani C, Gencer B, Collet TH, Rodondi N. Subclinical thyroid dysfunction and cardiovascular diseases: 2016 update. *Eur Heart J* 2018; 39(7): 503-7. <http://dx.doi.org/10.1093/eurheartj/ehx050> PMID: 28329380
- [4] Kravets I. Hyperthyroidism: Diagnosis and Treatment. *Am Fam Physician* 2016; 93(5): 363-70. PMID: 26926973
- [5] Wilson SA, Stem LA, Bruehlman RD. Hypothyroidism: Diagnosis and treatment. *Am Fam Physician* 2021; 103(10): 605-13. PMID: 33983002
- [6] Diab N, Daya NR, Juraschek SP, et al. Prevalence and risk factors of thyroid dysfunction in older adults in the community. *Sci Rep* 2019; 9(1): 13156. <http://dx.doi.org/10.1038/s41598-019-49540-z> PMID: 31511587
- [7] Juárez-Cedillo T, Basurto-Acevedo L, Vega-García S, et al. Prevalence of thyroid dysfunction and its impact on cognition in older Mexican adults: (SADEM study). *J Endocrinol Invest* 2017; 40(9): 945-52.

- [8] Awad SAS, Ashraf EM, Khaled AS, et al. The epidemiology of thyroid diseases in the Arab world: A systematic review. *J Public Health Epidemiol* 2016; 8(2): 17-26. <http://dx.doi.org/10.5897/JPHE2015.0713>
- [9] Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: An updated guideline for reporting systematic reviews. *Int J Surg* 2021; 88: 105906. <http://dx.doi.org/10.1016/j.ijsu.2021.105906> PMID: 33789826
- [10] Imani EF, Aminorroaya A, Soheilipour F, et al. Sonographic and functional characteristics of thyroid nodules in a population of adult people in Isfahan. *Endokrynol Pol* 2010; 61(2): 188-91. PMID: 20464706
- [11] Barzegari E, Pasdar Y, Hamzeh B, Darabi M, Moradinazar M, Najafi F. Factors associated with thyroid dysfunction in hypothyroidism-endemic region of ravansar, 2017-2018: A cross-sectional analysis. *J Adv Medi Biomed Res* 2021; 29(134): 145-51. <http://dx.doi.org/10.30699/jamb.29.134.145>
- [12] Abbasi M, Eshraghi B, Nikdel M. Epidemiological status of thyroid eye disease in Central Iran. *J Curr Ophthalmol* 2022; 34(1): 106-11. http://dx.doi.org/10.4103/joco.joco_223_21 PMID: 35620368
- [13] Niafar M, Najafipour F, Bahrami A. Subclinical thyroid disorders in postmenopausal women of Iran. *J Clin Diagn Res* 2009; 3: 1853-8.
- [14] Rezaei M, Javadmoosavi SY, Mansouri B, Azadi NA, Mehrpour O, Nakhaee S. Thyroid dysfunction: how concentration of toxic and essential elements contribute to risk of hypothyroidism, hyperthyroidism, and thyroid cancer. *Environ Sci Pollut Res Int* 2019; 26(35): 35787-96. <http://dx.doi.org/10.1007/s11356-019-06632-7> PMID: 31701424
- [15] Mehran L, Amouzegar A, Rahimabad P, Tohidi M, Tahmasebinejad Z, Azizi F. Thyroid function and metabolic syndrome: a population-based thyroid study. *Horm Metab Res* 2017; 49(3): 192-200. <http://dx.doi.org/10.1055/s-0042-117279> PMID: 28351085
- [16] Aminorroaya A, Amini M, Hovsepian S. Prevalence of hyperthyroidism in Isfahan-Iran, in the year 2006, fifteen years after universal salt iodization: a community based study. *Acta Endocrinol (Bucur)* 2008; 4(3): 273-85. <http://dx.doi.org/10.4183/aeb.2008.273>
- [17] Rahmani F, Rahmani F, Niafar M, Ostadi A, Robai N, Abdollahi F, et al. Thyroid function tests and the serum levels of prolactin in women with suicide attempt. *International Journal of Medical Toxicology and Forensic Medicine* 2018; 8(1): 5-10.
- [18] Aydin Y, Besir FH, Erkan ME, et al. Spectrum and prevalence of nodular thyroid diseases detected by ultrasonography in the Western Black Sea region of Turkey. *Med Ultrason* 2014; 16(2): 100-6. <http://dx.doi.org/10.11152/mu.2013.2066.162.ya1fhh2> PMID: 24791840
- [19] Demirba MB, Ucak S, Demirba ZS. Evaluation of thyroid diseases in hospitalized geriatric patients: A retrospective review. *Acta Med Mediter* 2019; 35(2): 1145-50.
- [20] Kutluturk F, Yildirim B, Ozturk B, et al. Thyroid dysfunctions and sonographic characteristics in northern Turkey: a population-based study. *Ann Saudi Med* 2013; 33(3): 253-9. <http://dx.doi.org/10.5144/0256-4947.2013.253> PMID: 23793427
- [21] Hergenç G, Onat A, Albayrak S, Karabulut A, Türkmen S. TSH levels in Turkish adults: Prevalences and associations with serum lipids, coronary heart disease and metabolic syndrome. *Turk J Med Sci* 2005; 35(5): 297-304.
- [22] Çayan S, Kendirci M, Yaman O, et al. Prevalence of erectile dysfunction in men over 40 years of age in Turkey: Results from the Turkish society of andrology male sexual health study group. *Turk J Urol* 2017; 43(2): 122-9. <http://dx.doi.org/10.5152/tud.2017.24886> PMID: 28717533
- [23] Sarıtaş A, Sarıtaş PU, Kurnaz MM, Çelik A. Spectrum and prevalence of thyroid disorders in patients admitted to the anaesthesiology outpatient clinic for surgery. *Turk J Anaesthesiol*

- Reanim 2015; 43(4): 240-5.
<http://dx.doi.org/10.5152/TJAR.2015.03206> PMID: 27366505
- [24] Rana H, Mirah JA, Al-Shahrani N, Nouf A, Afrah A, Basma O. Incidence of thyroid diseases in female Saudi adults visiting a tertiary care hospital in Riyadh. *Epidemiology* 2017; 7(1): 286.
- [25] Al Sayed N, Al Waili K, Alawadi F, *et al.* Consensus clinical recommendations for the management of plasma lipid disorders in the Middle East. *Int J Cardiol* 2016; 225: 268-83.
<http://dx.doi.org/10.1016/j.ijcard.2016.09.081> PMID: 27741487
- [26] Al-Shammari MA, Abdel Wahab MM, AlShamlan NA, *et al.* Clinical, laboratory, and ultrasound related diagnoses of thyroid disorders: Using a family medicine center data to assess thyroiditis and thyroid nodules in the eastern province of Saudi Arabia. *J Prim Care Community Health* 2022; 13: 21501319221095345.
<http://dx.doi.org/10.1177/21501319221095345> PMID: 35465762
- [27] Alqahtani SAM. Prevalence and characteristics of thyroid abnormalities and its association with anemia in asir region of Saudi Arabia: A cross-sectional study. *Clin Pract* 2021; 11(3): 494-504.
<http://dx.doi.org/10.3390/clinpract11030065> PMID: 34449542
- [28] Alanazi SL, Abo El-Fetoh NM, Alenezi HM, Alanazi WL, Alanazi KF, Alruwaili AS, *et al.* Pattern of thyroid diseases in arar city, northern Saudi Arabia. *Egypt J Hosp Med* 2018; 70(10): 1834-41.
<http://dx.doi.org/10.12816/0044762>
- [29] Abdulrahman Ibrahim AM, Alotaibi AMD. Survey of awareness of thyroid disorders among the Riyadh population, Central Region of Saudi Arabia. *Egypt J Hosp Med* 2018; 72(2): 4039-44.
<http://dx.doi.org/10.21608/ejhm.2018.9095>
- [30] Al Eidan E, Ur Rahman S, Al Qahtani S, Al Farhan AI, Abdulmajeed I. Prevalence of subclinical hypothyroidism in adults visiting primary health-care setting in Riyadh. *J Community Hosp Intern Med Perspect* 2018; 8(1): 11-5.
<http://dx.doi.org/10.1080/20009666.2017.1422672> PMID: 29441159
- [31] Refaat B. Prevalence and characteristics of anemia associated with thyroid disorders in non-pregnant Saudi women during the childbearing age: A cross-sectional study. *Biomed J* 2015; 38(4): 307-16.
<http://dx.doi.org/10.4103/2319-4170.151032> PMID: 25673171
- [32] Fureeh A, Al-Ghamdi A, Alhuussaini J, Alzahrani M, Alzahrani T. Prevalence and risk factors of subclinical thyroid disorders in Albaha region, Saudi Arabia. *Epidemiology* 2019; 9(368): 2161-1165.
- [33] Qashqary M, Tobaiqy M, Al-Sutari MM, Mujallad A, Alsheikh I. Prevalence of suspected cases of hyperthyroidism in jeddah by using wayne's scoring index. *Cureus* 2020; 12(11): e11538.
<http://dx.doi.org/10.7759/cureus.11538> PMID: 33354482
- [34] Alghaithy AA, El Reweny A, Shaaban Y, Hamdy A. The role of TSH receptor antibody *versus* thyroid peroxidase and thyroglobulin antibodies in detecting immune thyroid diseases in Saudi Patients at Almadinah Almounawarah. *Life Sci J* 2013; 10: 2900-7.
- [35] Alyahya A, AlNaim A, AlBahr AW, Almansour F, Elshebiny A. Knowledge of thyroid disease manifestations and risk factors among residents of the eastern province, Saudi Arabia. *Cureus* 2021; 13(1): e13035.
<http://dx.doi.org/10.7759/cureus.13035> PMID: 33665056
- [36] SJ Aljabri K, M Alnasser I, Facharatz B, *et al.* The frequency of hypothyroidism in Saudi community-based hospital: A retrospective single centre study. *Trends in Diabetes and Metabolism* 2019; 2(1): 1-4.
<http://dx.doi.org/10.15761/TDM.1000107>
- [37] Suhail N, Alsel BTA, Batool S. Prevalence and association of thyroid dysfunction with anemia/body iron status among northern Border Saudi population. *Int J Med Res Health Sci* 2020; 9(3): 1-7.
- [38] Abu-Helalah M, Alshraideh HA, Al-Sarayreh SA, *et al.* A cross-sectional study to assess the prevalence of adult thyroid dysfunction disorders in Jordan. *Thyroid* 2019; 29(8): 1052-9.
<http://dx.doi.org/10.1089/thy.2018.0579> PMID: 31146635
- [39] Zaman B, Rasool SO, Sabri SM, *et al.* Prevalence of thyroid dysfunctions in a large, unselected population in Duhok city, Iraqi Kurdistan: A cross-sectional study. *Journal of Biological Research Bollettino della Società Italiana di Biologia Sperimentale* 2021; 94(2)
<http://dx.doi.org/10.4081/jbr.2021.10067>
- [40] Al-Msari AMA, AL-Duleimy AHS, Lateef IA. Evaluation of patients with thyroid diseases in Baquba City according to thyroid function tests. *Diyala Journal of Medicine* 2014; 7(1): 76-9.
- [41] Salih SM, Kamel WA, Abbas MT, Abass KS. Prevalence of hyperthyroidism and hypothyroidism and its correlation with serum antithyroglobulin among patients in Kirkuk-Iraq . *J Adv Pharm Educ Res* 2021; 11(2): 57-60.
<http://dx.doi.org/10.51847/kWVVD06AagO>
- [42] Hamasaeed P, Hussain S, Ashraf S. Evaluation of Thyroid stimulating hormone and thyroid hormone concentrations in females with hypothyroidism and hyperthyroidism. *Rafidain Journal of Science* 2019; 28(4): 1-7.
<http://dx.doi.org/10.33899/rjs.2019.163290>
- [43] Shakor FN. Assessment of the clinical concern about the non-malignant thyroid disorders in Sulaimaniyah city-Kurdistan Region-Iraq. *Kufa Journal for Nursing Sciences* 2019; 9(2)
- [44] Tahir N, Najim H, Nsaif A. Prevalence of overt and subclinical thyroid dysfunction among Iraqi population in Baghdad city. *Iraqi J Community Med* 2020; 33(1): 20-4.
http://dx.doi.org/10.4103/IRJCM.IRJCM_3_20
- [45] Deraz H, Shawk N, Abednabi ALS, Ali El-Feitouri G. Study of Thyroid Profiles in the Elderly. *Zagazig Univ Med J* 2019; 25(5): 648-56.
<http://dx.doi.org/10.21608/zumj.2019.7008.10100>
- [46] Mahmoud AH. Thyroid dysfunction in elderly patients and its clinical presentation. *Thyroid* 2005; 14: 17.
- [47] Rashad NM, Samir GM. Prevalence, risks, and comorbidity of thyroid dysfunction: a cross-sectional epidemiological study. *Egypt J Intern Med* 2019; 31(4): 635-41.
http://dx.doi.org/10.4103/ejim.ejim_22_19
- [48] Okati-Aliabad H, Ansari-Moghaddam A, Kargar S, Mohammadi M. Prevalence of hypertension and pre-hypertension in the Middle East region: a systematic review & meta-analysis. *J Hum Hypertens* 2022; 36(9): 794-804.
<http://dx.doi.org/10.1038/s41371-021-00647-9> PMID: 35031669
- [49] Okati-Aliabad H, Ansari-Moghaddam A, Kargar S, Jabbari N. Prevalence of obesity and overweight among adults in the middle east countries from 2000 to 2020: A systematic review and meta-analysis. *Journal of Obesity* 2022; 2022(1): 8074837.
- [50] Garmendia Madariaga A, Santos Palacios S, Guillén-Grima F, Galofré JC. The incidence and prevalence of thyroid dysfunction in Europe: a meta-analysis. *J Clin Endocrinol Metab* 2014; 99(3): 923-31.
<http://dx.doi.org/10.1210/jc.2013-2409> PMID: 24423323
- [51] Mendes D, Alves C, Silverio N, Batel Marques F. Prevalence of undiagnosed hypothyroidism in Europe: a systematic review and meta-analysis. *Eur Thyroid J* 2019; 8(3): 130-43.
<http://dx.doi.org/10.1159/000499751> PMID: 31259155
- [52] Thavaraputta S, Dennis JA, Laoveeravat P, Nugent K, Rivas AM. Hypothyroidism and its association with sleep apnea among adults in the United States: NHANES 2007-2008. *J Clin Endocrinol Metab* 2019; 104(11): 4990-7.
<http://dx.doi.org/10.1210/jc.2019-01132> PMID: 31305928
- [53] Okati-Aliabad H, Ansari-Moghadam A, Mohammadi M, Kargar S, Shahrazi-Sanavi F. The prevalence of anxiety and depression and its association with coping strategies, supportive care needs, and social support among women with breast cancer. *Support Care Cancer* 2022; 30(1): 703-10.
<http://dx.doi.org/10.1007/s00520-021-06477-2> PMID: 34365523
- [54] Sajjadi-Jazi SM, Sharifi F, Varmaghani M, Meybodi HA, Farzadfar F, Larijani B. Epidemiology of hyperthyroidism in Iran: a systematic review and meta-analysis. *J Diabetes Metab Disord* 2018; 17(2): 345-55.
<http://dx.doi.org/10.1007/s40200-018-0367-1> PMID: 30918870